

On a covering problem in the hypercube

Lale Özkahya

email: ozkahya@illinoisalumni.org

Brendon Stanton

The Citadel

Department of Mathematics

Charleston, SC 29409 USA

email: bstanton@citadel.edu

October 4, 2011

Abstract

In this paper, we address a particular variation of the Turán problem for the hypercube. Alon, Krech and Szabó (2007) asked “In an n -dimensional hypercube, Q_n , and for $\ell < d < n$, what is the size of a smallest set, S , of Q_ℓ ’s so that every Q_d contains at least one member of S ?” Likewise, they asked a similar Ramsey type question: “What is the largest number of colors that we can use to color the copies of Q_ℓ in Q_n such that each Q_d contains a Q_ℓ of each color?” We give upper and lower bounds for each of these questions and provide constructions of the set S above for some specific cases.

1 Introduction

For graphs Q and P , let $\text{ex}(Q, P)$ denote the *generalized Turán number*, i.e., the maximum number of edges in a P -free subgraph of Q . The n -dimensional hypercube, Q_n , is the graph whose vertex set is $\{0, 1\}^n$ and whose edge set is the set of pairs that differ in exactly one coordinate. For a graph G , we use $n(G)$ and $e(G)$ to denote the number of vertices and the number of edges of G , respectively.

In 1984, Erdős [8] conjectured that

$$\lim_{n \rightarrow \infty} \frac{\text{ex}(Q_n, C_4)}{e(Q_n)} = \frac{1}{2}.$$

Note that this limit exists, because the function above is non-increasing for n and bounded. The best upper bound $\text{ex}(Q_n, C_4)/e(Q_n) \leq 0.62256$ was obtained by Thomason and Wagner [16] by slightly improving the bound 0.62284 given by Chung [4]. Brass, Harborth and Nienborg [3] showed that the lower bound is $\frac{1}{2}(1 + 1/\sqrt{n})$, when $n = 4^r$ for integer r , and $\frac{1}{2}(1 + 0.9/\sqrt{n})$, when $n \geq 9$.

Erdős [8] also asked whether $o(e(Q_n))$ edges in a subgraph of Q_n would be sufficient for the existence of a cycle C_{2k} for $k > 2$. The value of $\text{ex}(Q_n, C_6)/e(Q_n)$ is between $1/3$ and 0.3941 given by Conder [6] and Lu [11], respectively. On the other hand, nothing is known for the cycle of length 10. Except C_{10} , the question of Erdős is answered positively by showing that $\text{ex}(Q_n, C_{2k}) = o(e(Q_n))$ for $k \geq 4$ in [4], [7] and [10].

A generalization of Erdős' conjecture above is the problem of determining $\text{ex}(Q_n, Q_d)$ for $d \geq 3$. As for $d = 2$, the exact value of $\text{ex}(Q_n, Q_3)$ is still not known. The best lower bound for $\text{ex}(Q_n, Q_3)/e(Q_n)$ has been $1 - (5/8)^{0.25} \approx 0.11086$ due to Graham, Harary, Livingston and Stout [14] until recently Offner [12] improved it to 0.1165 . The best upper bound is $\text{ex}(Q_n, Q_3)/e(Q_n) \leq 0.25$ due to Alon, Krech and Szabó [1]. They also gave the best bounds for $\text{ex}(Q_n, Q_d)$, $d \geq 4$, as

$$\Omega\left(\frac{\log d}{d2^d}\right) = 1 - \frac{\text{ex}(Q_n, Q_d)}{e(Q_n)} \leq \begin{cases} \frac{4}{(d+1)^2} & \text{if } d \text{ is odd,} \\ \frac{4}{d(d+2)} & \text{if } d \text{ is even.} \end{cases} \quad (1)$$

These Turán problems are also asked when vertices are removed instead of edges and most of these problems are also still open. In a very recent paper, Bollobás, Leader and Malvenuto [2] discuss open problems on the vertex-version and their relation to Turán problems on hypergraphs.

Here, we present results on a similar dual version of the hypercube Turán problem that is asked by Alon, Krech and Szabó in [1]. Let \mathcal{H}_n^i denote the collection of Q_i 's in Q_n for $1 \leq i \leq n-1$. Call a subset of \mathcal{H}_n^ℓ a (d, ℓ) -covering set if each member of \mathcal{H}_n^d contains some member of this set, i.e., \mathcal{H}_n^d is covered by this set. A smallest (d, ℓ) -covering set is called *optimal*. Alon, Krech and Szabó [1] asked what the size of the optimal (d, ℓ) -covering set of Q_n is for fixed $\ell < d$. Call this function $f^{(\ell)}(n, d)$. Determining this function when $\ell = 1$ is equivalent to the determination of $\text{ex}(Q_n, Q_d)$, since $\text{ex}(Q_n, Q_d) + f^{(1)}(n, d) = e(Q_n)$ and the best bounds for $f^{(1)}(n, d)$ are given in [1] as (1). In [1], also the Ramsey version of this problem is asked as follows. A coloring of \mathcal{H}_n^ℓ is d, ℓ -polychromatic if all colors appear on each copy Q_d 's. Let $pc^{(\ell)}(n, d)$ be the largest number of colors for which there exists a d, ℓ -polychromatic coloring of \mathcal{H}_n^ℓ .

Let $c^{(\ell)}(n, d)$ be the ratio of $f^{(\ell)}(n, d)$ to the size of \mathcal{H}_n^ℓ , i.e.,

$$c^{(\ell)}(n, d) = \frac{f^{(\ell)}(n, d)}{2^{n-\ell} \binom{n}{\ell}}. \quad (2)$$

One can observe that

$$c^{(\ell)}(n, d) \leq \frac{1}{pc^{(\ell)}(n, d)}, \quad (3)$$

since any color class used in a d, ℓ -polychromatic coloring is a (d, ℓ) -covering set of Q_n . Note that the following limits exist, since $c^{(\ell)}(n, d)$ is non-decreasing, $pc^{(\ell)}(n, d)$ is non-increasing and both are bounded.

$$c_d^{(\ell)} = \lim_{n \rightarrow \infty} c^{(\ell)}(n, d), \quad p_d^{(\ell)} = \lim_{n \rightarrow \infty} pc^{(\ell)}(n, d).$$

In Section 2, we obtain bounds on the polychromatic number.

Theorem 1. *For integers $n > d > \ell$, let $0 < r \leq \ell + 1$ such that $r = d + 1 \pmod{\ell + 1}$. Then*

$$e^{\ell+1} \left(\frac{d+1}{\ell+1} \right)^{\ell+1} \geq \binom{d+1}{\ell+1} \geq p_d^{(\ell)} \geq \left\lceil \frac{d+1}{\ell+1} \right\rceil^r \left\lfloor \frac{d+1}{\ell+1} \right\rfloor^{\ell+1-r} \approx \left(\frac{d+1}{\ell+1} \right)^{\ell+1}.$$

In Section 3, we present the following bounds on $c_d^{(\ell)}$ and $c^{(\ell)}(n, d)$.

Theorem 2. *For integers $n > d > \ell$ and $r = d - \ell \pmod{\ell + 1}$,*

$$\left(2^{d-\ell} \binom{d}{\ell} \right)^{-1} \leq c_d^{(\ell)} \leq \left\lceil \frac{d+1}{\ell+1} \right\rceil^{-r} \left\lfloor \frac{d+1}{\ell+1} \right\rfloor^{-(\ell+1-r)}.$$

The determination of the exact values of $p_d^{(\ell)}$ and $c_d^{(\ell)}$ remains open. The lower and upper bounds on $c^{(\ell)}(n, d)$ provided in Theorem 2 and Theorem 3, respectively, are a constant factor of each other when d and ℓ have a bounded difference from n .

Theorem 3. *Let $n - d$ and $n - \ell$ be fixed finite integers, where $d > \ell$. Then, for sufficiently large n ,*

$$c^{(\ell)}(n, d) \leq \left\lceil \frac{r \log(n - \ell)}{\log\left(\frac{r^r}{r^r - r!}\right)} \right\rceil \frac{1 + o(1)}{2^{d-\ell} \binom{d}{\ell}},$$

where $r = n - d$.

Finally, we show an exact result for $c^{(\ell)}(n, d)$ when $d = n - 1$.

Theorem 4. *For integers $n - 1 > \ell$,*

$$c^{(\ell)}(n, n - 1) = \frac{\left\lceil \frac{2n}{n-\ell} \right\rceil}{2^{n-\ell} \binom{n}{\ell}}.$$

In our proofs, we make use of the following terminology. The collection of i -subsets of $[n] = \{1, \dots, n\}$, $1 \leq i \leq n$, is denoted by $\binom{[n]}{i}$. For an edge $e \in E(Q_n)$, $\text{star}(e)$ denotes the coordinate that is different at endpoints of e . The set of coordinates whose values are 0 (or 1, resp.) at both endpoints of e are denoted by $\text{zero}(e)$ (or $\text{one}(e)$, resp.). For a subcube $F \subset Q_n$, $\text{star}(F) := \cup_{e \subseteq E(F)} \text{star}(e)$, $\text{one}(F) := \cap_{e \subseteq E(F)} \text{one}(e)$ and $\text{zero}(F) := \cap_{e \subseteq E(F)} \text{zero}(e)$. Note that E_1 covers E_2 for $E_1 \in \mathcal{H}_n^\ell$ and $E_2 \in \mathcal{H}_n^d$ ($d > \ell$) if and only if $\text{zero}(E_2) \subset \text{zero}(E_1)$ and $\text{one}(E_2) \subset \text{one}(E_1)$.

Definition 5. For any $Q \in \mathcal{H}_n^\ell$ and $\text{star}(Q)$ with coordinates $s_1 < s_2 < \dots < s_\ell$, we define an $(\ell + 1)$ -tuple $w(Q) = (w_1, w_2, \dots, w_{\ell+1})$ as

- $w_1 = |\{x \in \text{one}(Q) : x < s_1\}|$,
- $w_j = |\{x \in \text{one}(Q) : s_{j-1} < x < s_j\}|$, for $2 \leq j \leq \ell$,
- $w_{\ell+1} = |\{x \in \text{one}(Q) : x > s_\ell\}|$.

2 Polychromatic Coloring of Subcubes

Proof of Theorem 1. *The lower bound:*

For any $Q \in \mathcal{H}_n^\ell$ with $w(Q) = (w_1, w_2, \dots, w_{\ell+1})$, we define the color of each $Q \in \mathcal{H}_n^\ell$ as the $(\ell + 1)$ -tuple $c(Q) = (c_1, \dots, c_{\ell+1})$ such that

$$\begin{aligned} c_i &= w_i \pmod{k} & \text{if } 1 \leq i \leq r \text{ and} \\ c_i &= w_i \pmod{k'} & \text{if } r + 1 \leq i \leq \ell + 1, \end{aligned} \tag{4}$$

where $k = \lceil (d + 1)/(\ell + 1) \rceil$ and $k' = \lfloor (d + 1)/(\ell + 1) \rfloor$. We show that this coloring is d, ℓ -polychromatic.

Let $C \in \mathcal{H}_n^d$, where $\text{star}(C)$ consists of the coordinates $a_1 < a_2 < \dots < a_d$. We choose a color $(c_1, \dots, c_{\ell+1})$ arbitrarily and show that C contains a copy of Q_ℓ , call it Q , with this color.

Since Q must be a subgraph of C , $\text{zero}(C) \subset \text{zero}(Q)$ and $\text{one}(C) \subset \text{one}(Q)$. We define $\text{star}(Q) = \{s_1, \dots, s_\ell\}$ such that

$$s_i = \begin{cases} a_{ik} & \text{if } 1 \leq i \leq r, \\ a_{rk+(i-r)k'} & \text{if } r + 1 \leq i \leq \ell. \end{cases}$$

We include the remaining $d - \ell$ positions of $\text{star}(C)$ to $\text{one}(Q)$ or $\text{zero}(Q)$ such that $w(Q) = (w_1, w_2, \dots, w_{\ell+1})$ satisfies (4). This is possible since by the definition of r , we have $d - \ell = r(k - 1) + (\ell + 1 - r)(k' - 1)$.

The upper bound:

Since $pc^{(\ell)}(n, d)$ is a non-increasing function of n , we provide an upper bound for this function when n is sufficiently large which is also an upper bound for $p_d^{(\ell)}$.

For a subset S of $[n]$, we define $\text{cube}(S)$ as the subcube Q of Q_n such that $\text{star}(Q) = S$ and $\text{zero}(Q) = [n] \setminus S$. Let \mathcal{G} be a subfamily of \mathcal{H}_n^d such that $\mathcal{G} = \{\text{cube}(S) : S \in \binom{[n]}{d}\}$. We define a coloring of the members of \mathcal{G} as follows.

Consider a d, ℓ -polychromatic coloring of \mathcal{H}_n^ℓ using p colors, call this coloring P . Fix an arbitrary ordering of the copies of Q_ℓ 's in Q_d . We define a coloring of \mathcal{H}_n^d such that the color of a copy of Q_d is the list of colors of each Q_ℓ under P in this fixed order. By using this coloring on the members of \mathcal{G} , we obtain a coloring of \mathcal{G} using $p^{(d)} 2^{d-\ell}$ colors.

Now, consider the auxiliary d -uniform hypergraph \mathcal{G}' whose vertex set is the set of coordinates $[n]$ and whose edge set is defined as the collection of $\text{star}(E)$'s for each E in \mathcal{G} , i.e., \mathcal{G}' is a complete d -uniform hypergraph on the vertex set $[n]$. Also we define a coloring of the edges of \mathcal{G}' by using the colors on the corresponding members of \mathcal{G} as described above. Ramsey's theorem on hypergraphs implies that there is a sufficiently large value of n such that there exists a complete monochromatic subgraph on $d^2 + d - 1$ vertices in any edge coloring of \mathcal{G}' with $p^{\binom{d}{\ell} 2^{d-\ell}}$ colors. Let $K \subset [n]$ be the vertex set of a monochromatic complete subgraph of \mathcal{G}' on $d^2 + d - 1$ vertices. We define S as the collection of $i d^{\text{th}}$ coordinates in K , $1 \leq i \leq d$, so that there are at least $d - 1$ coordinates between elements of S .

Claim 6. *If Q is a copy of Q_ℓ in $\text{cube}(S)$, then the color of Q under P depends only on $w(Q)$.*

Proof. Let E_1 and E_2 be two different copies of Q_ℓ in $\text{cube}(S)$ such that $w(E_1) = w(E_2)$ according to Definition 5. There exists a subset $S' \subset K$ with $|S'| = d$ such that

- $(\text{one}(E_2) \cup \text{star}(E_2)) \subset S'$, i.e., E_2 is contained in $\text{cube}(S')$ and
- the restriction of E_2 on S' gives the same vector as the restriction of E_1 on S .

Clearly, one can find S' that satisfies the first condition. It is also possible that S' fulfills the second condition, since we can remove or add up to $d - 1$ coordinates from K between consecutive coordinates of ones and stars in E_2 to define S' . This implies that the colors of E_1 and E_2 are the same under P , since $\text{cube}(S)$ and $\text{cube}(S')$ have the same colors. \square

Hence, the number of colors used in any d, ℓ -polychromatic coloring of \mathcal{H}_n^ℓ is at most the number of possible vectors $w(Q)$ for any $Q \in \mathcal{H}_n^\ell$. The number of possible $(\ell + 1)$ -tuples $w(Q)$ for any $Q \in \mathcal{G}$ is given by the number of partitions of at most $d - \ell$ ones into $\ell + 1$ parts and therefore it is at most $\binom{d+1}{\ell+1}$. \square

3 The Covering Problem

Proof of Theorem 2. Note that a trivial lower bound on $f^{(\ell)}(n, d)$ is given by the ratio of $|\mathcal{H}_n^d|$ to the exact number of Q_d 's that a single Q_ℓ covers in Q_n . Thus, by (2), for all n ,

$$c^{(\ell)}(n, d) \geq \left\lceil \frac{2^{n-d} \binom{n}{d}}{\binom{n-\ell}{n-d}} \right\rceil \cdot \frac{1}{2^{n-\ell} \binom{n}{\ell}}. \quad (5)$$

By using the equality $\binom{n}{d} \binom{d}{d-\ell} = \binom{n}{\ell} \binom{n-\ell}{d-\ell}$, we are done.

The upper bound is implied together by (3) and Theorem 1. \square

We define a $(0, 1)$ -labelling of a set as an assignment of labels 0 or 1 to its elements.

Observation 7. *Since any subcube $Q \subset Q_n$ is defined by $\text{zero}(Q)$ and $\text{one}(Q)$, a (d, ℓ) -covering set of Q_n can be defined as a collection of $(0, 1)$ -labellings of sets chosen from $\binom{[n]}{n-\ell}$ such that any $(0, 1)$ -labelling of sets in $\binom{[n]}{n-d}$ is contained in at least one of the labelled $(n - \ell)$ -sets.*

When providing constructions for the upper bounds in Theorems 3 and 4, we provide constructions for the equivalent covering problem in Observation 7.

Proof of Theorem 3.

We construct a (d, ℓ) -covering of Q_n by providing a construction for the equivalent problem as stated in Observation 7. In the following, we describe this construction in two steps. First, we choose the $(n - \ell)$ -subsets of $[n]$ to label and then, we describe an efficient way to $(0, 1)$ -labellings these sets.

Step 1: We make use of the following well-known result on the general covering problem. An (n, k, t) -covering is defined as a collection of k -subsets of n elements such that every t -set is contained in at least one k -set. Let $C(n, k, t)$ be the minimum number of k -sets in an (n, k, t) -covering. Rödl proved the following result by also settling a long-standing conjecture of Erdős and Hanani [9]. For any fixed integers k and t with $2 \leq t < k < n$,

$$\lim_{n \rightarrow \infty} \frac{C(n, k, t)}{\binom{n}{t} / \binom{k}{t}} = 1. \quad (6)$$

By our assumption, $n - d$ and $n - \ell$ are fixed integers where $n - d < n - \ell$. By (6), there exists a $(n, n - \ell, n - d)$ -covering \mathcal{F} for sufficiently large n such that $|\mathcal{F}| = (1 + o(1)) \binom{n}{n-d} / \binom{n-\ell}{n-d}$.

Step 2: We obtain a collection of $(0, 1)$ -labellings for each edge $e \in \mathcal{F}$ so that all $(0, 1)$ -labellings of $(n - d)$ -subsets of e are covered. The union of these $(0, 1)$ -labellings is a covering set.

An r -cut of an r -uniform hypergraph is obtained by partitioning its vertex set into r parts and taking all edges that meet every part in exactly one vertex. An r -cut cover of a hypergraph is a collection of r -cuts such that each edge is in at least one of the cuts. An upper bound on the minimum size of an r -cut cover is shown by Cioabă, Kündgen, Timmons and Vysotsky in [5] using a probabilistic proof.

Theorem 8 ([5]). *For every r , an r -uniform complete hypergraph on n vertices can be covered with $\lceil c \log n \rceil$ r -cuts if*

$$c > \frac{-r}{\log \left(\frac{r^r - r!}{r^r} \right)}.$$

For a fixed edge e of \mathcal{F} , let \mathcal{G}_e be the complete $(n - d)$ -uniform hypergraph on the vertex set of e . Let $C = \lceil c \log(n - \ell) \rceil$ be the size of a minimum $(n - d)$ -cut cover of \mathcal{G}_e as given by Theorem 8. We obtain a collection of $(0, 1)$ -labellings of e by labelling each cut in this cover such that the vertices in each part are labelled identically with 0 or 1. Thus, the total number of $(0, 1)$ -labellings of e is $2^{n-d} C$. (If some labelling of an edge is used more than once, then we count this labelling only once.) Finally, we use similarly labellings for each edge of \mathcal{F} in the covering set. This yields that

$$c^{(\ell)}(n, d) \leq \frac{1}{2^{n-\ell} \binom{n}{\ell}} (C(1 + o(1)) 2^{n-d} \frac{\binom{n}{n-d}}{\binom{n-\ell}{n-d}}) = C(1 + o(1)) \frac{1}{2^{d-\ell} \binom{d}{\ell}},$$

where the last equality is obtained by using the relation $\binom{n}{d}\binom{d}{d-\ell} = \binom{n}{\ell}\binom{n-\ell}{d-\ell}$. □

Proof of Theorem 4.

The lower bound follows from (5).

For the upper bound, we construct a collection of $(0, 1)$ -labellings of sets chosen from $\binom{[n]}{n-\ell}$, where singletons in $[n]$ have both 0 and 1 in some labelling. Let $k = \lceil n/(n-\ell) \rceil$. We choose a partition $[n] = (P_1, \dots, P_k)$ such that $|P_i| = n-\ell$ for $i < k$. Let $P \in \binom{[n]}{n-\ell}$ such that $P_k \subset P$. In the covering set, we include two labellings of each of P_1, \dots, P_{k-1}, P , where all labels are the same, either 0 or 1. □

References

- [1] N. Alon, A. Krech and T. Szabó. Turán's theorem in the hypercube. *SIAM J. Discrete Math.*, 21: 66–72, 2007.
- [2] B. Bollobás, I. Leader and C. Malvenuto. Daisies and other Turán Problems. *Combin., Probab. and Comp.*, 20: 743–747, 2011.
- [3] P. Brass, H. Harborth and H. Nienborg. On the Maximum number of edges in a C_4 -free subgraph of Q_n . *J. Graph Theory*, 19: 17–23, 1995.
- [4] F. Chung. Subgraphs of a hypercube containing no small even cycles. *J. Graph Theory*, 16: 273–286, 1992.
- [5] S. M. Cioabă, A. Kündgen, C. M. Timmons and V. V. Vysotsky. Covering complete r -graphs with spanning complete r -partite r -graphs. *Combin., Probab. and Comput.*, 20: 519–527, 2011.
- [6] M. Conder. Hexagon-free subgraphs of hypercubes. *J. Graph Theory*, 17: 477–479, 1993.
- [7] D. Conlon. An extremal theorem in the hypercube. *Electron. J. Combin.*, 17: R111, 2010.
- [8] P. Erdős. On some problems in graph theory combinatorial analysis and combinatorial number theory. *Graph Theory and Combinatorics*, 1–17, 1984.
- [9] P. Erdős and H. Hanani. On a limit theorem in combinatorial analysis. *Publ. Math. Debrecen*, 10: 10–13, 1963.
- [10] Z. Füredi, L. Özkahya. On even-cycle-free subgraphs of the hypercube. *J. of Combin. Theo., Ser. A*, 118: 1816–1819, 2011.

- [11] Linyuan Lu. Hexagon-free subgraphs in hypercube Q_n . Private communication.
- [12] D. Offner. Some Turán type results on the hypercube. *Discrete Math.*, 102: 2905–2912, 2011.
- [13] D. Offner. Polychromatic colorings of the hypercube. *SIAM J. Discrete Math.*, 22: 450–454, 2008.
- [14] N. Graham, F. Harary, M. Livingston and Q. Stout. Subcube fault tolerance in hypercubes. *Inform. and Comput.*, 102: 280–314, 1993.
- [15] V. Rödl. On a packing and covering problem. *European J. Combin.*, 6: 69–78, 1985.
- [16] A. Thomason and P. Wagner. Bounding the size of square-free subgraphs of the hypercube. *Discrete Math.*, 309: 1730–1735, 2009.